

## Joslin Diabetes Center and Joslin Clinic Guideline for Detection and Management of Diabetes in Pregnancy

9/10/2010: Revised 06-15-11

The Joslin Guideline for Detection and Management of Diabetes in Pregnancy is designed to assist internal medicine specialists, endocrinologists and obstetricians in individualizing the care of and setting goals for women with pre-existing diabetes who are pregnant or planning pregnancy. It is also a guide for managing women who are at risk for or who develop Gestational Diabetes Mellitus (GDM). This Guideline is not intended to replace sound medical judgment or clinical decision-making. Clinical judgment determines the need for adaptation in all patient care situations; more or less stringent interventions may be necessary.

The objectives of the Joslin Guideline for Detection and Management of Diabetes in Pregnancy are to support clinical practice and to influence clinical behaviors in order to improve clinical outcomes and assure that patient expectations are reasonable and informed. Guidelines are developed and approved through the Clinical Oversight Committee that reports to the Chief Medical Officer of the Joslin Diabetes Center. The Guideline is established after careful review of current evidence, medical literature and sound clinical practice. This Guideline will be reviewed periodically and modified as clinical practice evolves and medical evidence suggests.

### SCREENING FOR GESTATIONAL DIABETES MELLITUS

See *Screening Strategy to Detect GDM* Algorithm

### PRECONCEPTION CARE

#### Pre-existing type 1 or type 2 diabetes

<b>Glucose Goals Prior to Conception</b>	<ul style="list-style-type: none"> <li>• Fasting and pre-meal blood glucose: <b>Plasma*</b> 80-110 mg/dl</li> <li>• 1 hour postprandial blood glucose: 100-155 mg/dl</li> <li>• A1C &lt; 7%; and as close to normal as possible without resulting severe hypoglycemia</li> <li>• Avoid severe hypoglycemia</li> </ul>
<b>Counseling</b>	<ul style="list-style-type: none"> <li>• Educate women of childbearing age about the importance of near normal blood glucose control prior to conception</li> <li>• Refer to a maternal fetal-medicine and/or endocrinologist/diabetes specialist for counseling, assessment of maternal and fetal risk and guidance in achieving management goals. This includes all women who are planning pregnancy and women who are not planning pregnancy but are using inadequate contraception and have an A1C greater than 7%.</li> <li>• Assess diabetes self-management, including meal plan, insulin care and use, activity program, medication schedule, self-monitoring of blood glucose (SMBG), treatment for hypo- and hyperglycemia, and sick day management, using diabetes educators (DE) as appropriate. Review maternal and fetal health issues.</li> <li>• Begin a multivitamin with 400 mcgs folic acid to supplement average daily intake of 400 mcgs for a total daily intake of 800 mcg to 1 mg of folic acid to decrease the risk of neural tube defects. Patients with a prior pregnancy affected with a neural tube defect should take folic acid 4 mgs daily. Check a B12 level in patients consuming more than 1 mg folic acid, as high dose folic acid may mask B12 deficiency.</li> <li>• Strongly advise smoking and alcohol cessation</li> <li>• Refer overweight and obese women with and without known diabetes or polycystic ovary syndrome (PCOS) for medical nutrition therapy with a goal of 5-10% weight loss based on Institute of Medicine (IOM) 2009 recommendation.</li> </ul>
<b>Medical Assessment</b>	<ul style="list-style-type: none"> <li>• Medical and obstetrical history: including comprehensive review of diabetes history and management</li> <li>• Eye evaluation: dilated comprehensive eye exam and pregnancy clearance by an ophthalmologist; should also include a discussion about the risk of developing and/or the progression of diabetic retinopathy during pregnancy</li> <li>• Renal evaluation: spot urine microalbumin and serum creatinine; protein/creatinine ratio if spot urine microalbumin &gt;300 mcg/mg</li> <li>• Thyroid evaluation: TSH level</li> <li>• GYN evaluation: pelvic exam, Pap smear up to date</li> <li>• Cardiac evaluation: if ≥ 35 years of age with one or more additional risk factors (hypertension, smoking, family history of CAD, hypercholesterolemia, microalbuminuria or nephropathy) - recommend one or more of the following: EKG, echocardiogram, exercise tolerance test</li> </ul>
<b>Diabetes Medications</b>	<ul style="list-style-type: none"> <li>• Discontinue oral antihyperglycemic therapy; start insulin. An exception is metformin, which may be continued during the first trimester in patients with PCOS or type 2 diabetes, and anovulatory infertility. At the first prenatal visit the patient should begin increasing doses of insulin as necessary to control blood glucose while metformin is tapered off or discontinued. Metformin should not be used beyond the first trimester or in lieu of insulin until randomized controlled studies evaluating safety and efficacy have been completed. <ul style="list-style-type: none"> <li>○ Metformin crosses the placenta and achieves therapeutic levels in the fetus. Presently, there are no long term randomized controlled safety data in infants whose mother's were treated with metformin in pregnancy.</li> </ul> </li> <li>• Oral medications have not been adequately studied for the treatment of preexisting type 2 diabetes in pregnancy.</li> <li>• The rapid-acting insulin analogs lispro and aspart lower postprandial blood glucose and decrease the risk of nocturnal hypoglycemia and may be useful therapeutic agents. Patients on lispro and aspart prior to conception may continue them during pregnancy. Patients on regular insulin may be switched to lispro or aspart if 1-hour postprandial blood glucose levels are above target and the patient is also experiencing pre-meal or nocturnal hypoglycemia.</li> </ul>

	<ul style="list-style-type: none"> <li>• Glargine, a long-acting insulin analog, is not recommended in women who are planning a pregnancy or who are currently pregnant. There is no information on its safety in pregnancy. A specific concern in the pregnant population is related to the 6 to 8 fold increased IGF-1 receptor affinity and mitogenic potency compared with human insulin.</li> <li>• There is no information on the safety of using the two new insulin analogs, glulisine and detemir, in pregnancy. We cannot recommend their use at this time except in clinical trials.</li> <li>• The rapid-acting insulins, lispro or aspart may be delivered either through multiple daily injections (MDI) or an insulin pump.</li> <li>• There is inadequate safety information about the use of exenatide and liraglutide in pregnancy. They should therefore not be used in pregnancy.</li> </ul>
<b>Other Medications</b>	<p><b>Hypertension and/or microalbuminuria management:</b></p> <ul style="list-style-type: none"> <li>• ACE-inhibitors must be stopped before pregnancy due to the increased risk of birth defects in the 1st trimester and the risk of fetal injury or demise with 2nd or 3rd trimester use. ARBs should be stopped before conception because safety data for 1st trimester use is limited. <ul style="list-style-type: none"> <li>○ The non-dihydropyridine calcium channel blocker diltiazem in extended release forms may be a useful substitute for ACE-Is and ARBs.</li> </ul> </li> </ul> <p><b>Diabetic nephropathy/chronic renal disease management:</b></p> <ul style="list-style-type: none"> <li>• Sometimes the benefits of preconception use of ACE-inhibitors for renal protection may outweigh the mildly increased risk of birth defects. In this case, ACE-inhibitors should be stopped as soon as pregnancy is diagnosed in the first trimester.</li> </ul> <p><b>Hyperlipidemia management:</b></p> <ul style="list-style-type: none"> <li>• Stop all cholesterol-lowering agents before conception, including statins</li> <li>• Hypertriglyceridemia—omega 3 fatty acids may be started or continued in pregnancy</li> </ul>

## DIABETES MANAGEMENT DURING PREGNANCY

<b>Self Monitoring of Blood Glucose and Urine Ketones</b>	<p><b><u>Pre-existing diabetes and GDM</u></b></p> <ul style="list-style-type: none"> <li>• For gestational diabetes, check glucose levels 4 times/day: before breakfast and 1 hour post-meals</li> <li>• For pre-existing diabetes, check glucose levels pre-meals and 1 hour post-meal</li> <li>• Nocturnal monitoring (~3 AM) may be necessary on an intermittent basis</li> <li>• Check fasting urine ketones daily</li> </ul>																																							
<b>Treatment Goals</b>	<table border="0"> <thead> <tr> <th></th> <th style="text-align: center;"><b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b></th> <th style="text-align: center;"><b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b></th> </tr> </thead> <tbody> <tr> <td>• Fasting and pre-meal plasma glucose</td> <td style="text-align: center;">60-99 mg/dl</td> <td style="text-align: center;">60-79 mg/dl</td> </tr> <tr> <td>• 1-hour post-meal or peak post-prandial plasma glucose</td> <td style="text-align: center;">100-129 mg/dl</td> <td style="text-align: center;">90-109 mg/dl</td> </tr> <tr> <td>• Urine ketones</td> <td style="text-align: center;">negative</td> <td style="text-align: center;">negative</td> </tr> <tr> <td>• Normalization of hemoglobin A1C to &lt; 6% if possible without resulting severe hypoglycemia</td> <td></td> <td></td> </tr> <tr> <td>• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)</td> <td></td> <td></td> </tr> <tr> <td>• Avoidance of severe hypoglycemia (episode in which patient experiences coma, seizure or suspected seizure, or impairment sufficient to require the assistance of another person). Blood glucose goals must be relaxed for patients with hypoglycemia unawareness or recurrent hypoglycemia.</td> <td></td> <td></td> </tr> </tbody> </table> <hr style="border-top: 1px dashed black;"/> <table border="0"> <thead> <tr> <th></th> <th style="text-align: center;"><b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b></th> <th style="text-align: center;"><b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b></th> </tr> </thead> <tbody> <tr> <td>• Fasting and premeal blood glucose</td> <td style="text-align: center;">60-95 mg/dl</td> <td style="text-align: center;">60-79 mg/dl</td> </tr> <tr> <td>• 1-hour post meal or peak post prandial</td> <td style="text-align: center;">100-129 mg/dl</td> <td style="text-align: center;">90-109 mg/dl</td> </tr> <tr> <td>• Urine ketones</td> <td style="text-align: center;">negative</td> <td style="text-align: center;">negative</td> </tr> <tr> <td>• Initiate insulin therapy if above levels are not maintained</td> <td></td> <td></td> </tr> <tr> <td>• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)</td> <td></td> <td></td> </tr> </tbody> </table>		<b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b>	<b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b>	• Fasting and pre-meal plasma glucose	60-99 mg/dl	60-79 mg/dl	• 1-hour post-meal or peak post-prandial plasma glucose	100-129 mg/dl	90-109 mg/dl	• Urine ketones	negative	negative	• Normalization of hemoglobin A1C to < 6% if possible without resulting severe hypoglycemia			• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)			• Avoidance of severe hypoglycemia (episode in which patient experiences coma, seizure or suspected seizure, or impairment sufficient to require the assistance of another person). Blood glucose goals must be relaxed for patients with hypoglycemia unawareness or recurrent hypoglycemia.				<b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b>	<b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b>	• Fasting and premeal blood glucose	60-95 mg/dl	60-79 mg/dl	• 1-hour post meal or peak post prandial	100-129 mg/dl	90-109 mg/dl	• Urine ketones	negative	negative	• Initiate insulin therapy if above levels are not maintained			• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)		
	<b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b>	<b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b>																																						
• Fasting and pre-meal plasma glucose	60-99 mg/dl	60-79 mg/dl																																						
• 1-hour post-meal or peak post-prandial plasma glucose	100-129 mg/dl	90-109 mg/dl																																						
• Urine ketones	negative	negative																																						
• Normalization of hemoglobin A1C to < 6% if possible without resulting severe hypoglycemia																																								
• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)																																								
• Avoidance of severe hypoglycemia (episode in which patient experiences coma, seizure or suspected seizure, or impairment sufficient to require the assistance of another person). Blood glucose goals must be relaxed for patients with hypoglycemia unawareness or recurrent hypoglycemia.																																								
	<b>Plasma glucose*</b> <b>Hadlock AC &lt; 70<sup>th</sup> percentile</b>	<b>Plasma glucose*</b> <b>Hadlock AC ≥ 70<sup>th</sup> percentile</b>																																						
• Fasting and premeal blood glucose	60-95 mg/dl	60-79 mg/dl																																						
• 1-hour post meal or peak post prandial	100-129 mg/dl	90-109 mg/dl																																						
• Urine ketones	negative	negative																																						
• Initiate insulin therapy if above levels are not maintained																																								
• Use standard hypoglycemia treatment for blood glucose less than 60 mg/dl (15 grams of carbohydrate – recheck in 15 minutes; repeat with 15 grams of carbohydrate if blood glucose is still less than 60 mg/dl)																																								
<b>Diabetes Monitoring and Visits</b>	<p><b><u>Pre-existing Diabetes</u></b></p> <ul style="list-style-type: none"> <li>• Medical visits (endocrinologist preferred) every 1-4 weeks, with additional phone contact as needed, depending on level of self-management skills and stability of blood glucose control. At each visit, review SMBG and urine ketone results, measure blood pressure, measure urine protein and ketones by dipstick</li> <li>• Check A1C level every 4-8 weeks</li> <li>• Education using a diabetes educator (DE), preferably a Certified Diabetes Educator (CDE), as needed; medical nutrition therapy (MNT) by registered dietitian (RD)</li> <li>• Ophthalmology exam early in first trimester; follow-up depending on findings of this exam</li> <li>• Consider providing mental health counseling to assist women and / or their partners cope with the psychological and relationship changes that may result from pregnancy.</li> </ul>																																							

	<p><b><u>Gestational Diabetes</u></b></p> <ul style="list-style-type: none"> <li>• Medical visits (endocrinologist preferred) every 1-4 weeks, with additional phone contact as needed, depending on level of self-management skills and stability of blood glucose control. At each visit, review SMBG and urine ketone results, measure blood pressure, measure urine protein and ketones by dipstick.</li> <li>• If newly diagnosed with gestational diabetes, patient should be started on insulin, not metformin, if medication is required.</li> <li>• Education using DE (preferably a CDE) as needed for review of SMBG to increase adherence; MNT by registered dietitian (RD)</li> <li>• Insulin is preferred over glyburide in GDM as studies comparing glyburide to insulin were not powered to evaluate neonatal outcomes. There was a trend to greater infant birth weights when mothers were treated with glyburide compared to insulin.</li> <li>• When insulin is not an option, glyburide may be used.</li> </ul>
<b>Diabetes Medications</b>	<p><b><u>Preexisting Diabetes</u></b></p> <ul style="list-style-type: none"> <li>• The only diabetes medication currently used throughout pregnancy is insulin (see <b>Preconception Care</b>).</li> </ul>
<b>Hypertension Management</b>	<ul style="list-style-type: none"> <li>• Maintaining blood pressure in non-pregnant patients at <math>\leq 130/80</math> decreases end organ damage.</li> <li>• Target blood pressure is 110-129 systolic and 65-79 diastolic in women with chronic hypertension during pregnancy. Antihypertensives are initiated in pregnant patients with known or suspected chronic hypertension if blood pressure is <math>\geq 130/80</math> three times during pregnancy.</li> <li>• Pre-eclampsia needs special treatment; therefore, these guidelines and treatment strategies do not apply to pre-eclampsia for which other treatment options are preferred, or to gestational hypertension when high blood pressure exposure is limited</li> <li>• Antihypertensives that are used during pregnancy are: <ul style="list-style-type: none"> <li>➢ Alpha methyldopa (category B)</li> <li>➢ Beta-blockers (acebutolol, sotalol – category B; betaxolol, bisoprolol, labetalol, levatol, metoprolol, nadolol, timolol – category C; atenolol – category D – should not be used as it may cause fetal growth restriction)</li> <li>➢ Calcium channel blockers (all category C) (The nondihydropyridine calcium channel blocker diltiazem in extended-release form may be preferred in patients with microalbuminuria or nephropathy.)</li> <li>➢ Hydralazine (category C)</li> </ul> </li> </ul>

\*Laboratory methods measure plasma glucose. Most glucose monitors approved for home provide readings equivalent to plasma values. Plasma glucose values are 10-15% higher than whole blood glucose values. It is important for people with diabetes to know whether their meters and strips record whole blood or plasma results.

## MEDICAL NUTRITION THERAPY (MNT)

*Recommendations are the same for pre-existing diabetes and GDM except where noted.*


<b>Counseling and Education</b>	<ul style="list-style-type: none"> <li>• All pregnant women should receive MNT counseling by a registered dietitian (RD), (CDE preferred)</li> <li>• All pregnant women should receive SMBG training by a DE (CDE preferred)</li> <li>• Daily food records and SMBG records are required to assess effectiveness of MNT</li> <li>• Carbohydrate counting skills are taught for either a consistent carb intake or a personalized insulin to carb ratio so the patient can adjust insulin based on carbohydrate intake</li> <li>• At least 3 encounters with a CDE are recommended: <ul style="list-style-type: none"> <li>○ Visit 1 (60 – 90 min individual or group visit with RD) for assessment and meal planning. This could include SMBG instruction if RD has received appropriate training.</li> <li>○ Visit 2 (30 – 45 min) with RD or RN 1 week after initial visit to assess and modify plan</li> <li>○ Visit 3 (15 – 45 min) with RD or RN in 1 – 3 weeks to further assess and modify plan, as needed.</li> </ul> </li> <li>• Additional visits every 2 – 3 weeks and prn with RD or RN until delivery, and one visit 6 – 8 weeks after delivery</li> </ul>			
<b>Calories</b>	WHO BMI range (kg/m <sup>2</sup> )	Energy Needs (kcal/kg) °Based on Pregravid kg	Total Weight Gain Range (lbs) <b>Twin</b>	Rates of weight Gain (lb/week) 2 <sup>nd</sup> and 3 <sup>rd</sup> trimesters
	Underweight <18.5	36-40	28-40	1.0 (1-3)
	Normal weight 18.5-24.9	30	25-35      37-54	1.0 (0.8-1)
	Overweight 25.0-29	24	15-25      31-50	0.6 (0.5-0.7)
	Obese $\geq 30.0$	**	11-20      25-42	0.5 (0.4-0.6)
<p>For singleton pregnancy, add an additional 340 kcal/day to calculated needs in 2<sup>nd</sup> trimester and 452 kcal/day in 3<sup>rd</sup> trimester, or additional calories consistent with target weight gain. For twin pregnancy, add an additional 500 kcals to calculated needs after 1<sup>st</sup> trimester. For multiple pregnancies, add 500 kcal in the 1<sup>st</sup> trimester. (**Provisional guidelines for twin pregnancies per Institute of Medicine (IOM)).</p> <p>* Insufficient information was available to develop a provisional guideline for underweight women with multiple fetuses.  ** Insufficient information to address energy needs (kcal/kg) in the obese category</p>				

<b>Distribution of Calories</b>	<ul style="list-style-type: none"> <li>Individualize distribution of calories based on usual intake, preferences and medication regimen <ul style="list-style-type: none"> <li>6 – 8 small meals/snacks is recommended. Smaller frequent meals decrease postprandial hyperglycemia</li> </ul> </li> </ul> <p>Weight should be monitored at each visit; track patient's weight gain on prenatal weight gain chart</p>		
<b>Carbohydrate</b>	<b><u>GDM</u></b>	<b><u>Pre-Existing Diabetes</u></b>	
	40 – 45% total calories*	40% – 55% total calories	
	<b>Breakfast</b>	15 – 30 grams* +	Individualized as per usual intake and BG levels
	<b>Other meals</b>	45 grams lunch and dinner	Individualized as per usual intake and BG levels
	<b>HS snack</b>	15 – 30 grams carbohydrate	15 – 30 grams carbohydrate
	*Pregnant women should consume a minimum of 175 grams of carbohydrate per day + May be increased if insulin is added		
<b>Fiber</b>	<ul style="list-style-type: none"> <li>Calculate 14 grams of fiber per 1000 kcals per day (25-30 grams/day) based on provider assessment</li> </ul>		
<b>Protein</b>	<ul style="list-style-type: none"> <li>Calculate 1.1 grams of protein per kg per day, based on provider assessment</li> </ul>		
<b>Fat</b>	<ul style="list-style-type: none"> <li><u>Pre-existing diabetes</u>: 30 – 40% of total calories, with &lt;10% total calories from saturated fat</li> <li><u>GDM</u>: 30-40% total calories with &lt;10% total calories from saturated fat</li> <li>Encourage use of monounsaturated and polyunsaturated fats instead of saturated fats</li> </ul>		
<b>Nutritive and Non-nutritive Sweeteners</b>	<ul style="list-style-type: none"> <li>The safety of non-nutritive sweeteners has not been established.</li> </ul>		
<b>Vitamin/ Mineral Supplements</b>	<p>Prenatal multivitamin and mineral supplement including:</p> <ul style="list-style-type: none"> <li>Iron (27 mg/day)</li> <li>Folic acid 400 mcgs to supplement average daily dietary intake of 400 mcgs for a total daily intake of 800 mcgs to 1 mg daily to decrease risk of neural tube defects (begin 400 mcg prior to conception)</li> <li>Additional calcium supplementation may be needed to meet daily requirement of 1000 mg per day (1300 mg per day if under age 19yrs). Begin prior to conception.</li> <li>Vitamin D 600 IUs/day.</li> </ul>		
<b>Caffeine</b>	<ul style="list-style-type: none"> <li>Limit to &lt;200 mg per day. Excess caffeine consumption during pregnancy may increase the risk of miscarriage.</li> </ul>		
<b>Physical Activity</b>	<ul style="list-style-type: none"> <li>Regular physical activity is recommended after clearance by provider <ul style="list-style-type: none"> <li>Benefits include reducing insulin resistance, postprandial hyperglycemia and excessive weight gain</li> </ul> </li> <li>Hypoglycemia is more likely with prolonged exercise (&gt;60 minutes)</li> <li>Encourage activity after meals to reduce postprandial hyperglycemia</li> </ul>		
<b>Alcohol and Tobacco Use</b>	<ul style="list-style-type: none"> <li>Alcohol and tobacco use should be avoided during pregnancy.</li> </ul>		

## POST-PARTUM CARE

- Breastfeeding is encouraged in patients with pre-existing or gestational diabetes
- Enalapril and captopril may be used to treat hypertension and albuminuria in nursing mothers of full-term infants
- Appointments with the following specialists should be completed 6-8 weeks post-partum: ophthalmology, RD or RN and endocrinology.
- For women who develop GDM:
  - A 2-hour 75 g OGTT should be checked at 6 weeks to evaluate for persistent diabetes
    - Normal: fasting glucose level <100mg/dl
    - Impaired fasting glucose: fasting glucose level 100-125mg/dl
    - Diabetes: fasting glucose level ≥126mg/dl
    - Impaired glucose tolerance: 2 hr OGTT value 140-199mg/dl
    - Diabetes: 2 hr OGTT value ≥ 200mg/dl
  - Counsel women with GDM on the role of lifestyle management and weight loss to reduce the risk of future type 2 DM (of note: approximately 50% of women with GDM will develop overt type 2 diabetes in the next 7 to 10 years)
  - Review nutrition guidelines and establish exercise goals. For women with BMI greater than 25 (this may be lower in Asians) target a 5-7% weight loss from the preconception weight.
- Discuss family planning/contraceptive issues. Depo-Provera and progestin-only oral contraceptives are less preferred in patients who have had gestational diabetes, as they can accelerate the development of type 2 diabetes. In patients with pre-existing diabetes, Depo-Provera may worsen glycemic control. The intrauterine device (IUD) is preferred in monogamous partnerships because it is a metabolically neutral and highly effective form of contraception.
- Assist women with gestational diabetes with the transfer of care back to the primary care physician for longer term diabetes screening (including yearly fasting glucose, 1 year post partum and every 3 years afterwards 75 gram 2hour OGTT), risk reduction and for lifestyle management.

**Gestational Diabetes Mellitus**  
 Screening Strategy to Detect GDM  
 Risk assessment should be done at first prenatal visit



\*\*Plasma values, based on International Association of Diabetes and Pregnancy Study Groups Consensus Panel (IADPSG)

## Glossary

**Hadlock AC:** formula to identify macrosomia established by Hadlock et al

**ACE Inhibitor:** angiotensin converting enzyme inhibitor

**ARBs:** angiotensin receptor blockers

**BMI:** body mass index

**CAD:** coronary artery disease

**CDE:** Certified Diabetes Educator

**DE:** diabetes educator; nurse or dietitian with advanced education in diabetes management

**GDM:** gestational diabetes mellitus

**IFG:** impaired fasting glucose

**IGF:** receptor: Insulin-like growth factor receptor

**IOM:** Institute of Medicine

**MDI:** multiple daily injections

**MNT:** Medical Nutrition Therapy

**OGTT:** oral glucose tolerance test

**PCOS:** polycystic ovarian syndrome

**SMBG:** self-monitoring of blood glucose

**TSH:** thyroid stimulating hormone

**WHO:** World Health Organization

## References:

1. American Dietetic Association. Diabetes Mellitus (DM) Type 1 and Type 2 Evidence-based Nutrition Practice Guidelines for Adults. <http://www.adaevidencelibrary.com/topic.cfm?cat=3253>. (Accessed August 25, 2010).
2. American College of Obstetrics and Gynecology: Practice Bulletin. *Clinical Management Guidelines for Obstetricians-Gynecologists*. No. 30, September 2001.
3. Briggs GG, Freeman RK, Yaffe SJ (eds). *Drugs in Pregnancy and Lactation*. Vol 8. Philadelphia, PA. Lippincott, Williams and Wilkins, 2008, p. 239-244, 613-620.
4. Brown FM, Goldfine AB. Diabetes and Pregnancy. In Kahn CR, et al, editor, *Joslin's Diabetes Mellitus*, 14<sup>th</sup> edition. Philadelphia, PA. Lippincott, Williams & Wilkins, 2005, p.1036-1047.
5. Glueck CJ et al. Metformin throughout pregnancy reduces the development of gestational diabetes in women with polycystic ovarian syndrome. *Fertil Steril* 2002; 77:520-5.
6. Jovanovic L. Metabolic and immunologic effects of insulin lispro in gestational diabetes. *Diabetes Care* 1999; 22:1422-1427.
7. Kahn, HS, Curtis, KM, Marchbanks, PA. Effects of injectable or implantable progestin-only contraceptives on insulin-glucose metabolism and diabetes risk. *Diabetes Care* 2003; 26:216-225.
8. Kitzmiller J, Jovanovich L, Brown F, Coustan D, Reader, D. Eds. *Managing Preexisting Diabetes and Pregnancy: Technical Reviews and Consensus Recommendations for Care*. Alexandria, VA, American Diabetes Association, 2008.
9. Kjos, SL. Contraception and the risk of type 2 diabetes mellitus in Latina women with prior gestational diabetes mellitus. *JAMA* 1998; 280:539-543.
10. Krolewski AS, Kosinski EJ et al. Magnitude and determinants of coronary artery disease in juvenile-onset insulin-dependent diabetes mellitus. *Am J Cardiol* 1987; 59:750-755.
11. Kurtzhals P et al. Correlations of receptor binding and metabolic and mitogenic potencies of insulin analogs designed for clinical use. *Diabetes* 2000; 49:999-1005.
12. Miller E, Hare JW, Cloherty et al. Elevated maternal hemoglobin A1C in early pregnancy and major congenital anomalies in infants of diabetic mothers. *N Engl J Med* 1981; 304:1331-1334.
13. Report of the National High Blood Pressure Education Program Working Group on High Blood Pressure in Pregnancy. *Am J Obstet Gynecol* 2000; 183(1): S1-S22.
14. Boyd E, Metzger TA, Buchanan DR, Coustan, de Leiva A, Dunger DB, Hadden DR, Hod M, Kitzmiller JL, Kjos SL, Oats JN, Pettitt DJ, Sacks DA, Zoupas C. Summary and recommendations of the Fifth International Workshop-Conference on Gestational Diabetes Mellitus. *Diabetes Care* 2007; 30:S251-S260.
15. Kitzmiller JL, Block JM, Brown FM, Catalano PM, Conway DL, Coustan DR, Gunderson E, Herman WH, Hoffman LD, Inturrisi M, Jovanovic LB, Kjos SI, Knopp RH, Montoro MN, Ogata ES, Paramsothy P, Reader DM, Rosenn BM, Thomas AM, Kirkman MS. Managing preexisting diabetes for pregnancy. *Diabetes Care* 2008; 31:1060-1079.
16. Rowan JA, Hague WM, Gao W, Battin MR, Moore MP; MiG Trial Investigators. Metformin versus insulin for the treatment of gestational diabetes. *N Engl J Med* 2008; 358:2003-15. Erratum in: *N Engl J Med* 2008; 359:106.
17. HAPO Study Cooperative Research Group, Metzger BE, Lowe LP, Dyer AR, Trimble ER, Chaovarindr U, Coustan DR, Hadden DR, McCance DR, Hod M, McIntyre HD, Oats JJ, Persson B, Rogers MS, Sacks DA. Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med* 2008; 358:1991-2002.

18. Jensen DM, Korsholm L, Ovesen P, Beck-Nielsen H, Moelsted-Pedersen L, Westergaard JG, Moeller M, Damm P. Peri-conceptional A1C and risk of serious adverse pregnancy outcome in 933 women with type 1 diabetes. *Diabetes Care* 2009; 32:1046-1048.
19. Hadlock FP, Deter RL, Harrist RB, Park SK. Fetal abdominal circumference as a predictor of menstrual age. *AJR* 1982; 139:367-370.
20. Vahratian A, Barber JS, Lawrence JM, Catherine Kim C. Family-planning practices among women with diabetes and overweight and obese women in the 2002 National Survey for Family Growth. *Diabetes Care* 2009; 32:1026-1031.
21. Rasmussen KM, Yaktine AL, eds. Committee to reexamine IOM pregnancy weight guidelines. Weight gain during pregnancy: reexamining the guidelines. Institute of Medicine; National Research Council, 2009. [http://www.nap.edu/catalog.php?record\\_id=12584](http://www.nap.edu/catalog.php?record_id=12584) (accessed August 10, 2010).
22. Hillhouse JH, Neiger R. Pregnancy and lactation. In: Gottschlich MM, Fuhrman MP, Hammond KA, Holcombe BJ, Seidner DL, eds. *The Science and Practice of Nutrition Support: A Case-Based Core Curriculum*. Dubuque, IA, Kendall/Hunt Publishing Company, 2001, p. 302-319.
23. Brown JE, Carlson M. Nutrition and multifetal pregnancy. *J Am Diet Assoc* 2000; 100:343-348.
24. American Diabetes Association. Nutrition recommendations and interventions for diabetes. *Diabetes Care* 2008; 31(suppl 1):S61-S78.
25. Kitzmiller, JL, Block JM, Brown FM. Managing preexisting diabetes for pregnancy. *Diabetes Care* 2008; 31:1060-1079.
26. A Report of the Panel on Macronutrients, Subcommittees on Upper Reference Levels of Nutrients and Interpretation and Uses of Dietary Reference Intakes, and the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. Food and Nutrition Board, Institute of Medicine of the National Academies, 2005. [http://www.nap.edu/openbook.php?record\\_id=10490&page=R1](http://www.nap.edu/openbook.php?record_id=10490&page=R1) (accessed August 10, 2010).
27. American Diabetes Association. Standards of medical care in diabetes – 2010. *Diabetes Care* 2010; 33 (suppl 1):S11-S61.
28. Weng X, Odouli R, Li DK. Maternal caffeine consumption during pregnancy and the risk of miscarriage: a prospective cohort study. *Am J Obstet Gynecol* 2008; 198:279.e1-279.e8.
29. Kjos SL. Contraception for Women with Diabetes. In *Diabetes in Women: Pathophysiology and Therapy*. Tsatsoulis A, Wyckoff J, Brown FM, Eds. New York, NY, Humana Press, 2009, p.167-180.
30. Rizzotto JM, Guisti J, Higgins L. Nutrition and Pregnancy. In *Diabetes in Women: Pathophysiology and Therapy*. Tsatsoulis A, Wyckoff J, Brown FM, Eds. New York, NY, Humana Press, 2009, p 255-272.
31. Shane-McWhorter L. *Complementary & Alternative Medicine (CAM) Supplement Use in People with Diabetes: A Clinician's Guide*. Alexandria, VA, American Diabetes Association, 2007, p. 60.
32. Halldorsson TI, Strom M, Petersen SB, Olsen SF. Intake of artificially sweetened soft drinks and risk of preterm delivery: A prospective cohort study in 59,334 Danish pregnant women. *Am J Clin Nutr* 2010; 92:626-33.
33. Yodyingyuad V, Bunyawong S. Effect of stevioside on growth and reproduction. *Hum Reprod* 1991; 6:158-165.
34. Saenphet K, Aritajat S, Saenphet S, Manosroi J, Manosroi A. Safety evaluation of aqueous extracts from Aegle marmelos and Stevia rebaudiana on reproduction of female rats. *Southeast Asian J Trop Med Public Health* 2006; 37 Suppl 3:203-5.
35. The Healthy Eating Pyramid. <http://www.hsph.harvard.edu/nutritionsource/what-should-you-eat/pyramid-full-story/index.html> (accessed August 31, 2010).
36. International Association of Diabetes and Pregnancy Study Groups Consensus Panel: International Association of Diabetes and Pregnancy Study Groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. *Diabetes Care* 2010; 33: 676-691
37. Schaefer-Graf UM, Kjos SL, Fauzan OH, Bühling KJ, Siebert G, Bühner C, Ladendorf B, Dudenhausen JW, Vetter K. A randomized trial evaluating a predominantly fetal growth-based strategy to guide management of gestational diabetes in caucasian women. *Diabetes Care* 2004; 27:297-302.
38. Vitamin D Fact Sheet: <http://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/>

### Pregnancy Guideline Task Force

Florence Brown, MD – Task Force Leader

Suzanne Ghiloni, RN, BSN CDE

Tracey O’Keeffe Lucier, RD, LDN, CDE

Jo-Anne Rizzotto, MEd, RD, CDE

Approved by Joslin Clinical Oversight Committee on 09/10/2010

### Joslin Clinical Oversight Committee

Om Ganda, MD – Chairperson

William Hsu, MD

Susan Sjostrom, JD

Richard Beaser, MD

Richard Jackson, MD

Kenneth Snow, MD

Elizabeth Blair, MS ANP-BC, CDE

Lori Laffel, MD, MPH

William Sullivan, MD

Amy Campbell, MS, RD, CDE

Medha Munshi, MD

Howard Wolpert, MD

Cathy Carver ANP-BC, CDE

Melinda Maryniuk, MEd, RD, CDE

John Zrebiec, LICSW

Jerry Cavallerano, OD, PhD

Jo-Anne Rizzotto, MEd, RD, CDE

Martin Abrahamson, MD (*ex officio*)

David Feinbloom, MD

Bijan Roshan, MD